

Dr. K. M. Ledding*
Plastic & Reconstructive surgeon
CORNWALL PROFESSIONAL BUILDING
#506 2125 11th Avenue
Regina, Saskatchewan
S4P 3X3

PHONE (306) 780- 3650

FAX (306) 780 3718

INSTRUCTIONS AFTER FILLER INJECTIONS

Please observe the following after treatment:

- Immediately after the treatment you may experience slight redness, swelling, tenderness and an itching sensation in the treated area. This is normal result of the injection. The inconvenience is temporary and generally disappears in 1-2 days.
- If irritation continues, or if other reactions occur, please contact our office immediately.
- The initial swelling and tenderness after lip treatment may last longer. Some patients experience these symptoms for about a week and the lips may look somewhat uneven during this time. This means that the result directly after the treatment should not be seen as the final result.
- Avoid touching the treated area for 6 hours following treatment.
- Until the initial swelling and redness have resolved, do not expose the treated area to intense heat or extreme cold.
- If you have previously suffered from facial cold sores, there is the risk that the treatment could contribute to an eruption of cold sores. Please inform Dr. Ledding if you have previously had cold sores, and Dr. Ledding will provide you with a prescription for an anti-viral medication to limit this risk.
- If you are using aspirin or any similar medication, be aware that these may increase the bruising and bleeding at the injection site.

If you have any questions or concerns about your treatment, please contact our office at (306) 780-3650.