

In Order to Minimize the Risk of Bruising Prior to Injections – Neuromodulator & Dermal Fillers

Avoid know blood thinners and drugs associated with bruising and bleeding for 10 days prior to your procedure. This includes the following:

1. **ASA Family:** Aspirin/Baby Aspirin, Coated Aspirin/ASA, Anacin/Excedrin, Fiorina, Robaxiasal.
2. **Arthritis Pills:** Anaprox/Naproxen, Feldene, Ibuprofen/Motrin/Advil, Voltaren/Diclofenac, Toradol, Aleve, Indocid or other DSAIDS (non-steroidal inflammatory drops).
3. **Medical Blood Thinners:** Plavix, Coumadin, Warfarin, Eliquis, Pradaxa, Xarelto.
4. **Herbals and Vitamins:** Supplements/multivitamins, Cod-liver Oil/Fish oil capsules, Garlic Pills, Green Tea, Vitamin E, Omega-3, 6 Fatty acids.

* If any of these medications are prescribed It is important that you check with your doctor first to insure it is safe to stop*

If you have any questions, please contact our office at 306-780-3650. We'd be happy to assist you.

Your Patient Care Team!



Ledding Plastic Surgery

CORNWALL PROFESSIONAL BUILDING
#314, 2125 11th Avenue, Regina, SK S4P 3X3
Phone: 306-780-3650 Fax: 306-780-3718

www.leddingplasticsurgery.com leddingoffice@sasktel.net